



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



By (2),  $\rho^2 \dot{\theta} = A$ , so that (1) gives

$$\ddot{\rho} - A^2 \rho^{-3} = -\mu \rho^{-2},$$

or

$$\dot{\rho} = \sqrt{-A^2 \rho^{-2} + 2\mu \rho^{-1} + B}.$$

The speed is

$$\sqrt{\rho^2 + \rho^2 \theta^2} = \sqrt{2\mu \rho^{-1} + B}.$$

Hence,

$$B = v^2 - 2\mu a^{-1}.$$

Hence,

$$\frac{d\rho}{d\theta} = \frac{\dot{\rho}}{\dot{\theta}} = \rho^2 A^{-1} \sqrt{v^2 - 2\mu a^{-1} + 2\mu \rho^{-1} - A^2 \rho^{-2}}$$

and therefore,

$$\theta = - \int_{a^{-1}}^{\rho^{-1}} A^{-1} \sqrt{v^2 - 2\mu a^{-1} + 2\mu z - A^2 z^2} dz;$$

that is,

$$\rho^{-1} = \mu A^{-2} - (\mu A^{-2} - a^{-1}) \cos \theta + \sqrt{v^2 A^{-2} - a^{-2}} \sin \theta,$$

which is the equation of the trajectory.

Writing this in the form,

$$\mu a^2 (1 - \cos \theta)^2 + \{2\mu a (1 - \cos \theta) (\cos \theta - a \rho^{-1}) - a^2 v^2 \sin^2 \theta\} A^2 + \{(\cos \theta - a \rho^{-1})^2 + \sin^2 \theta\} A^4 = 0,$$

and applying the condition for equal roots in  $A^2$ , we get for the envelope

$$\frac{4\mu a^2 v^2}{4\mu^2 - a^2 v^4} \frac{1}{\rho} = 1 - \frac{2\mu - a v^2}{2\mu + a v^2} \cos \theta,$$

the equation of an ellipse.

### 349 (Mechanics). Proposed by S. A. COREY, Albia, Iowa.

A 9-pound weight is attached to a string which passes over a smooth fixed pulley. The other end of the string is fastened to and supports a smooth pulley  $P_1$  of weight 1 pound over which passes a second string, one end attached to a 3-pound weight and the other end attached to and supporting another smooth pulley  $P_2$  of weight 1 pound. Over the pulley  $P_2$  passes a third string supporting weights 2 pounds and  $3\frac{1}{2}$  pounds.

If the system is acted upon by gravity alone show that the acceleration of the 9-pound weight,  $3\frac{1}{2}$ -pound weight, and pulley  $P_2$  are 0,  $\frac{1}{2}g$ , and  $\frac{1}{3}g$ , respectively.

Determine the motion of the weights when pulleys are not smooth, that is, when friction is present.

## II. SOLUTION BY WILLIAM HOOVER, Columbus, Ohio.

Calling the fixed pulley  $P$  and taking it as the origin of coördinates, we have for the  $x$ -coordinate of  $m_1$ ,  $x_1$ ; of  $P_1$ ,  $l_1 - x_1$ ; of  $m_3$ ,  $x_2 + l_1 - x_1$ ; of  $P_2$ ,  $l_1 + l_2 - (x_1 + x_2)$ ; of  $m_4$ ,  $x_3 + l_1 + l_2 - (x_1 + x_2)$ ; of  $m_6$ ,  $l_1 + l_2 + l_3 - (x_1 + x_2 + x_3)$ , where  $m_1$ ,  $m_3$ ,  $m_4$  and  $m_6$  are the masses of the 9-, 3-, 2-, and  $3\frac{1}{2}$ -pound weights, respectively.

Under the hypothesis that there is no friction, the equation of motion of system is

$$\begin{aligned} T = \frac{1}{2} m_1 \dot{x}_1^2 + \frac{1}{2} P_1 \dot{x}_1^2 + \frac{1}{2} m_3 (\dot{x}_2 - \dot{x}_1)^2 + \frac{1}{2} (\dot{x}_1 + \dot{x}_2)^2 + \frac{1}{2} m_4 (\dot{x}_3 - \dot{x}_1 - \dot{x}_2)^2 + \frac{1}{2} m_6 (\dot{x}_1 + \dot{x}_2 \\ + \dot{x}_3)^2 = m_1 g x_1 + P_1 g (l_1 - x_1) + m_3 g (x_2 + l_1 - x_1) + P_2 g \{l_1 + l_2 - (x_1 + x_2)\} \\ + m_4 g \{l_1 + l_2 + x_3 - (\dot{x}_1 + x_2)\} + C = V. \end{aligned} \quad (i)$$

Using Lagrange's equations of type

$$\frac{d}{dt} \frac{dT}{dx} - \frac{dT}{dx} = \frac{dV}{dx} \quad (ii)$$

there are

$$(a) \quad \frac{5}{8} \ddot{x}_1 + \frac{1}{8} \ddot{x}_2 + \frac{4}{3} \ddot{x}_3 = g(m_1 - P_1 - m_3 - P_2 - m_4 - m_6) = -\frac{4}{3}g,$$

$$(b) \quad \frac{1}{8} \ddot{x}_1 + \frac{3}{8} \ddot{x}_2 + \frac{4}{3} \ddot{x}_3 = g(m_3 - P_2 - m_4 - m_6) = -\frac{1}{8}g,$$

$$(c) \quad \frac{4}{3} \ddot{x}_1 + \frac{4}{3} \ddot{x}_2 + \frac{1}{3} \ddot{x}_3 = g(m_4 - m_6) = -\frac{4}{3}g,$$

$$\text{giving } \ddot{x}_1 = 0, \ddot{x}_2 = -\frac{g}{3}, \ddot{x}_3 = -\frac{g}{6}.$$